

INTRODUCTION TO ALGEBRAS OF PARTIAL TRIANGULATIONS

LAURENT DEMONET

ABSTRACT. The aim of this note is to give a gentle introduction to algebras of partial triangulations of marked surfaces, following the structure of a talk given during the 49th symposium on ring theory and representation theory, held in Osaka. This class of algebras, which always have finite rank, contains classical Jacobian algebras of triangulations of marked surfaces and Brauer graph algebras. We discuss representation theoretical properties and derived equivalences. All results are proven in [2], under slightly milder hypotheses.

1. THE ALGEBRA OF A PARTIAL TRIANGULATION

Let k be a unital ring and Σ be a connected compact oriented surface with or without boundary. We fix a non-empty finite set $\mathbb{M} \subset \Sigma$ of marked points (some of them may be on the boundary $\partial\Sigma$). For each $M \in \mathbb{M}$, we fix an invertible coefficient $\lambda_M \in k^\times$ and a multiplicity $m_M \in \mathbb{Z}_{>0}$. For simplicity, we suppose here that if Σ is a sphere then $\#\mathbb{M} \geq 5$ and if Σ is a disc then $\#\mathbb{M} \geq 3$.

Definition 1. An *arc* on (Σ, \mathbb{M}) is a continuous map $u : [0, 1] \rightarrow \Sigma$ satisfying:

- The restriction of u to $(0, 1)$ is an embedding into $\Sigma \setminus \mathbb{M}$;
- Extremities 0 and 1 are mapped to \mathbb{M} .

We consider arcs up to homotopy relative to their endpoints in $\Sigma \setminus \mathbb{M}$. Moreover, we exclude arcs that are homotopic to a marked point or to a boundary component, that is the closure of a connected component of $\partial\Sigma \setminus \mathbb{M}$.

In this note, for simplicity, we exclude arcs that are loops enclosing a unique marked point $M \in \mathbb{M}$ with $m_M \leq 2$. We do not make this restriction in [2].

Definition 2. We say that two arcs u and v are *compatible* if, up to homotopy, they are non-crossing. Then, a *partial triangulation* of (Σ, \mathbb{M}) is a set σ of arcs of (Σ, \mathbb{M}) that are pairwise compatible. If σ is a maximal partial triangulation and each connected component of $\partial\Sigma$ contains at least a marked point, σ is called a *triangulation*.


In order to define the algebra Δ_σ of a partial triangulation σ , we first need to construct a quiver Q_σ .

Definition 3. The quiver Q_σ has set of vertices σ , and arrows are winding in Σ between successive arcs around marked points counter-clockwisely. We call *bouncing path* of Q_σ any path of length 2 consisting of two arrows that are not successive around the same endpoint.

We give two examples to illustrate this definition:

The paper is in a final form and no version of it will be submitted for publication elsewhere.

Example 4. In the left example Σ is a disc with four marked points A, B, C and D (A and B are on the boundary). In the right one, Σ is a torus with two marked points M and N . The partial triangulations are depicted with thick lines.



Bouncing paths of the first example are ab, cd, de and ec . Bouncing paths of the second example are a^2 and b^2 .

To define the algebra Δ_σ , we need two more combinatorial concepts:


Definition 5. For each edge $u \in \sigma$ and endpoint M of u , we denote by $\omega_{u,M}$ the path of Q_σ going from u to u winding once around M if $M \notin \partial\Sigma$ and $\omega_{u,M} = 0$ if $M \in \partial\Sigma$.

Definition 6. A *small triangle* of σ is a triple (u, v, w) of arcs of σ together with three marked points M, N and P such that

- u is incident to M and N ;
- v is incident to N and P ;
- w is incident to P and M ;
- The union of u, v and w encloses clockwise (in the order u, v, w) a disc without marked point inside.

We now define the *algebra of the partial triangulation* σ by $\Delta_\sigma := kQ_\sigma/I_\sigma$ where I_σ is the ideal generated by the following relations:

- (1) For each $u \in \sigma$ that joins $M \in \mathbb{M}$ to $N \in \mathbb{M}$, we require $\lambda_M \omega_{u,M}^{m_M} = \lambda_N \omega_{u,N}^{m_N}$.
- (2) For each $u \in \sigma$ with endpoint M , we require $\omega_{u,M}^{m_M} q = 0$ for any arrow q of Q_σ .
- (3) Suppose that σ contains a small triangle M, N, P as in the following picture:



where α, β and γ are arrows, ω is the only possible simple path winding around M if $M \notin \partial\Sigma$ and $\omega = 0$ if $M \in \partial\Sigma$. We require the relation $\alpha\beta = \lambda_M(\omega\gamma)^{m_M-1}\omega$.

- (4) For any bouncing path $\alpha\beta$ that does not appear in case 3, we require $\alpha\beta = 0$.

Example 7. Relations for the left partial triangulation of Example 4 are:

$$a^{m_D} = (df)^{m_C} = (fd)^{m_C} = 0 \quad (1)$$

$$ab = 0 \quad (4)$$

$$cd = de = 0 \quad (3)$$

$$ec = \lambda_C (fd)^{m_C-1} f \quad (3)$$

and other relations are redundant.

Relations for the right partial triangulation are:

$$(ab)^{m_M} = (ba)^{m_M} \quad (1)$$

$$a^2 = b^2 = 0. \quad (4)$$

It turns out that algebras of partial triangulations are particularly well behaved. A first result about them is that this definition is compatible with the naive notion of a sub-partial triangulation:

Theorem 8. *Let $\tau \subset \sigma$. Then we have*

$$\Delta_\tau \cong e_\tau \Delta_\sigma e_\tau$$

where e_τ is the sum of the primitive idempotents of Δ_σ corresponding to the arcs of τ .

Notice that we have naturally $kQ_\tau \subset kQ_\sigma$. However, relations as defined in this note do not go through this inclusion. We have to take a more complicated variant of these relations, giving an isomorphic algebra, to obtain Theorem 8.

2. BRAUER GRAPH ALGEBRAS AND JACOBIAN ALGEBRAS OF SURFACES

We explain here that the class of algebras of partial triangulations contains two important classes of algebras.

Theorem 9. *If σ contains no small triangle, neither arc incident to $\partial\Sigma$, then Δ_σ is the Brauer graph algebra of σ considered as a ribbon graph. Moreover, any Brauer graph algebra is the algebra of a partial triangulation of a surface without boundary.*

We will not recall what a Brauer graph algebra is. For more details, see for example [4] or [5]. However, this definition is very close to the definition of the algebra of a partial triangulation and Theorem 9 is mostly straightforward.

Theorem 10. *If all m_M are invertible in k and σ is a triangulation, then Δ_σ is the Jacobian algebra of a quiver with potential (Q_σ, W_σ) . To define W_σ , consider the set E of small triangles T of σ up to rotation and for each of them denote by α_T , β_T and γ_T the three arrows as in the figure defining relation (3) earlier. Then, for each $M \in \mathbb{M}$ take arbitrarily an arc u_M incident to M . Then*

$$W_\sigma := \sum_{T \in E} \alpha_T \beta_T \gamma_T - \sum_{M \in \mathbb{M}} \frac{\lambda_M}{m_M} \omega_{u_M}^{m_M}$$

(as usual for potentials, terms are only well defined up to cyclic permutations).

Notice that if $m_M = 1$ for all $M \in \mathbb{M}$, we recover the usual Jacobian algebra of a surface as defined in [3]. Recall that the Jacobian algebra of (Q_σ, W_σ) is the quotient of the completed path algebra $\widehat{kQ_\sigma}$ by the cyclic derivatives of W_σ . So we get here an improvement as Δ_σ is directly defined from kQ_σ without completion.

3. ALGEBRAIC PROPERTIES OF Δ_σ

We have the following result about Δ_σ :

Theorem 11. *The k -algebra Δ_σ is a free k -module of rank*

$$\sum_{M \in \mathbb{M} \cap \partial\Sigma} \frac{d_M(d_M - 1)}{2} + \sum_{M \in \mathbb{M} \cap (\Sigma \setminus \partial\Sigma)} m_M d_M^2 + f$$

where, for $M \in \mathbb{M}$, d_M is the degree of M in the graph σ (without counting boundary components), and f is the number of arcs in σ with both endpoints on boundaries.

Example 12. The algebra of the left partial triangulation of Example 4 has rank $5 + 4m_C + m_D$, and the right one has rank $4m_M$.

More precisely, there is a k -basis of Δ_σ consisting of all strict and non-idempotent prefixes of all $\omega_{u,M}^{m_M}$, together with primitive idempotents and elements $\lambda_M \omega_{u,M}^{m_M} = \lambda_N \omega_{u,N}^{m_N}$.

The following property generalizes a known result for Brauer graph algebras and Jacobian algebras of surfaces without boundary:

Theorem 13. *If σ has no arc incident to the boundary, then Δ_σ is a symmetric k -algebra (i.e. $\text{Hom}_k(\Delta_\sigma, k) \cong \Delta_\sigma$ as Δ_σ -bimodules).*

4. REPRESENTATION TYPE OF Δ_σ


The next theorem permits to expect that the classification of Δ_σ -modules is possible:

Theorem 14. *If k is an algebraically closed field, then Δ_σ is of tame representation type.*

The proof of this result relies on a deformation theorem by Crawley-Boevey [1]. Indeed, the relations defining Δ_σ can be deformed to the relations of a Brauer graph algebra in a suited manner. Moreover, Brauer graph algebras are of tame representation type. Notice that unfortunately, these techniques do not permit to deduce directly the classification of Δ_σ -modules even though modules over Brauer graph algebras are known.

5. FLIP OF PARTIAL TRIANGULATIONS AND DERIVED EQUIVALENCES

Finally, we give a flip leading to derived equivalences. For an arc u in σ such that arcs marked by $+$ in the following diagrams are also in σ (in particular, they are not in $\partial\Sigma$), we define $\mu_u(\sigma)$ by replacing u by u^* defined in the following way:




We also define coefficients $\mu_u(\lambda)$. For a marked point $M \in \mathbb{M}$, $\mu_u(\lambda)_M = \lambda_M$ except in the following cases:

- In case (F1), if M is the topmost vertex of the figure then $\mu_u(\lambda)_M = -\lambda_M$.
- In case (F1), if M is the rightmost vertex of the figure then $\mu_u(\lambda)_M = (-1)^{m_M} \lambda_M$.
- In case (F2), if M is the unique marked point enclosed by u then $\mu_u(\lambda)_M = -\lambda_M$.

Then we get the following result:

Theorem 15. *There is a derived equivalence between Δ_σ and $\Delta_{\mu_u(\lambda)}^{\mu_u(\sigma)}$ where the second algebra is computed with respect to the coefficients $\mu_u(\lambda)$.*

Example 16. We consider the two following partial triangulations of a disc with three marked points, none of them are in $\partial\Sigma$:



They are related by a flip so the following algebras, obtained for $\lambda_M = \lambda_N = \lambda_P = 1$ and $m_M = m_N = m_P = m$ are derived equivalent:

$$k \left(\begin{array}{c} \bullet \\ \swarrow \quad \searrow \\ \bullet \quad \bullet \\ \nwarrow \quad \nearrow \\ \bullet \end{array} \right) \quad \text{and} \quad k \left(\begin{array}{c} \alpha \quad \bullet \quad \xrightarrow{\beta_2} \bullet \quad \xleftarrow{\gamma_2} \bullet \quad \delta \\ \quad \quad \quad \beta_1 \quad \quad \quad \gamma_1 \end{array} \right)$$

$$\frac{(x^2 - (yx)^{m-1}y, y^2)}{\left(\begin{array}{l} \beta_2\alpha - (\gamma_1\gamma_2\beta_2\beta_1)^{m-1}\gamma_1\gamma_2\beta_2, \\ \alpha\beta_1 - (\beta_1\gamma_1\gamma_2\beta_2)^{m-1}\beta_1\gamma_1\gamma_2, \\ \beta_1\beta_2 - \alpha^{m-1}, \gamma_1\delta, \delta\gamma_2, \gamma_2\gamma_1, \\ \delta^m - (\gamma_2\beta_2\beta_1\gamma_1)^m \end{array} \right)}.$$

REFERENCES

- [1] William Crawley-Boevey, *Tameness of biserial algebras*, Arch. Math. (Basel), **65**(5) (1995), 399–407.
- [2] Laurent Demonet, *Algebras of partial triangulations*, submitted to Adv. Math., arXiv: 1602.01592.
- [3] Daniel Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces*, Proc. Lond. Math. Soc. (3), **98**(3) (2009), 797–839.
- [4] Klaus W. Roggenkamp, *Biserial algebras and graphs*, in Algebras and modules, II (Geiranger, 1996), volume **24** of CMS Conf. Proc., Amer. Math. Soc., Providence, RI, (1998), 481–496.
- [5] Burkhard Wald and Josef Waschbsch, *Tame biserial algebras*, J. Algebra, **95**(2) (1985), 480–500.

GRADUATE SCHOOL OF MATHEMATICS
NAGOYA UNIVERSITY
FUROCHO, CHIKUSAKU
NAGOYA 464-8602 JAPAN

E-mail address: Laurent.Demonet@normalesup.org